

Enumeration of the fungi of
Ceylon Part II, containing remainder

110 THE REV. M. J. BERKELEY AND MR. C. E. BROOME

974. HYDNOCYSTIS THWAITESII, B. & Br. Ochracea, globosa, inflata,
denum plicata, ore parvo aperta, extus glaberrima; sporidiis ellipticis
(no. 1056).

On the ground. Peradeniya, Dec. 1868.

Sometimes seated on a tomentose stratum; $\frac{1}{2}$ -1 inch across;
sporidia .001 by .0005.

975. TUBER ZEYLANICUM, B. & Br. Minutum, luteum rugosum, centro
afflum (no. 1013).

On a piece of dead stick. Peradeniya, Dec.

Only a single specimen found, $\frac{1}{2}$ line across, unfortunately
without perfect fruit. No. 1093 is in the same condition.

SPIRÆRIACEI.

976. CORDICEPS MILITARIS, Fr. (No. 341.)

On dead pupæ. Nuwara Eliya; Central province, 1868.
Bright crimson, varying much in size.

977. C. BARNESEII, Thw. Stipe cylindrico velutino; clavula cylindrica,
apice sterili; conidiophoris ramosis stilboideis candidis, capitulis glo-
bosis (no. 1120 cum icono).

On larvae of *Melolontha*.
Conidia .0001 long.

978. C. SOBOLIFERA, B. Hook. *Lond. Journ.* 1843, p. 207 (*sub Sphæria*).
Torrubia sobolifera, Tul. *Sel. Fung. Carp.* iii. p. 10.

On larvae of some lamellicorn insect at the roots of coffee-trees. Bala-
godde,

culatas formantibus, margine crenato niveto tomentoso; ascis subclavatis;
sporidiis oblongis, extus hyalinis 3-septatis (no. 628).
On bark. Tropical forests south of the island. July 1868.
Sporidia .0005 long by .0025, brown, with a greenish endochrome.
No. 630, from the same locality, is a variety with scattered cups.

d. P. ALBO-RUPA, B. & Br. Hymenio late rufo, margine albo crenato; spori-
diis subfusiformibus 5-septatis, ascis clavatis (no. 69).
On bark. Peradeniya, Nov. 1867.

e. P. MAGNIPICA, B. & Br. Hymenio croceo-lateritio, cuticula elevata rigida
cineto, pulvorumento; sporidiis sigmoides multiseptatis, articulis constrictis
(no. 624).

On sticks. Tropical forests south of the island. July 1868.
Sporidia .004-.0045 long, .0005-.0007 wide, ultimately breaking up into
elliptic disks.

The magnificent sporidia remind one of those of *Lichenopsis sphaeroboloides*,
Schwein.

Bakley's M.J. and C.E. Broome 1868 or 1873?

Journal of the Linnean Society, London
of Hymenomycetes 14, 110-111 1875

111

979. *C. MYRMECOPHILA*, Cés. (sub *Hypoocrea*); Rab. (no. 1033). Ochroleuca; stipite filiformi tenacello; clavula ovoidea ad basin sterili, superne costata acutusecula, e peritheiorum summo ostiolo, gibberulosa (no. 1218 cum icono).

On dead ants.

980. *C. DIPTERIGENA*, B. & Br. Pallida, stipite cylindrico; capite globoso; ostiolis inconspicuis.

Sep. 1864.

About 1½ inch high.

Under no. 619 is some *Spermædia* on *Fimbristylis* from the south of the island, July 1863. The resulting *Cordiceps* has not yet been detected.

981. *EPICHLAE PULVINULUS*, B. & Br. Brevis, aurantiaca, pulvinata, ostiolis depressis (no. 1099).

On some species of *Panicum*. Peradeniya, Dec. 1868.

Forming little cushions, 1-2 lines long, confined to one side, but springing from the same sort of mycelium as *E. typhina*; asci linear, waved; sporidia with short articulation.

982. *E. CINerea*, B. & Br. Cylindrica, cinerea; sporis longis filiformibus (no. 504).

On *Eragrostis nutans*. March 6, 1868.

983. *HYPOCREA ARTOCREAS*, B. & Br. Hemisphaerica, late affixa, subtus tomentosa albida, supra plana jecorina rugosiuscula, primum pulverulenta (no. 110).

Apparently on petioles of some palms. Nov. 1868.

Sporidia not mature.

984. *H. DISCELLA*, B. & Br. Pallide rufa, orbicularis, applanata, supra rugosa punctata; sporidiis minoribus (no. 5 in part).

On dead wood.

Asci linear; sporidia .0001 long.

985. *H. DEPLANATA*, B. & Br. Cremoricolor, adnata, margine brevi albo byssideo; ostiolis impressis latioribus (no. 1100).

Central province, Dec. 1868.

Sporidia 16, globose, forming a necklace .0002 in diameter.

986. *H. CORTICOIDES*, B. & Br. Effusa, olivacea, margine brevi byssideo; peritheciis nunc immersis, nunc liberis collapsis; sporidiis ellipticis obtusis apiculatis uniscptatis (no. 645).

On bark. South of the island, July 1868.

Sporidia .0002-.00025 long.

987. *H. SACCHARINA*, B. & C. Journ. Linn. Soc. x. p. 376.