

tain dermatophytes, or ringworm (May) 1929.
gworm fungi. Biol. Rev. 4: 41-

The effects of hydrogen ion con-
amin C on the growth of fungi.
65, (Aug.) 1938.

An alkaline medium and procedure
tes in the presence of saprophytic
d. 30: 119-131, (Feb.) 1945.

ucts of pathogenic fungi. Biology
Cryptogamici et Phytopathologici,
Botanica Co., Waltham, Massa-

ference on dermatophytes. Medi-
onal Research Council, (Nov. 8)

NEW SPECIES OF TORRUBIELLA, HIRSU- TELLA AND GIBELLULA¹

E. B. MAINS

(WITH 2 FIGURES)

Torrubiella pulvinata sp. nov. (FIG. 1, A & B)

Mycelium album, corpus hospitis tegente et ad articulos floccos creante; perithecia partim immersa vel superficialia in pulvino mycelii, ovoidea, 450-600 × 220-270 μ, brunnea, a tenui tunica flavi mycelii tecta; asci cylindrici, 450 × 4.5-6 μ, deorsum attenuata, tenuibus membranibus, ad apices membranibus spissatis 2.5-3 μ; ascosporeae filiformes, 0.5-0.7 μ crassae, multiseptatae, cellis 4-8 μ longis; conidiophora e mycelio, sparsa vel caespitosa, 2-2.5 μ crassa, sursum capitulis phialidum; phialides anguste fusoido-ovoideae, sursum acuminatae, 8-12 × 2 μ; conidia hyalina, fusoido-ellipsoidea, 2-3 × 1.5-2 μ, catenulata.

Ex Opilionoideis, Waianuka, Oahu, Hawaii. Dec. 9, 1945, D. P. Rogers (2090).

Mycelium white, covering the body of the host and developing tufts on the legs specially at the joints; perithecia partly embedded or superficial on a pulvinate mass of mycelium, ovoid, 450-600 × 220-270 μ, brown, covered by a thin layer of yellow mycelium; asci cylindric up to 450 μ long, 4.5-6 μ wide, narrowing toward the base, the wall thin, thickened at the apex, 2.5-3 μ; ascospores filiform, 0.5-0.7 μ thick, multiseptate, the cells 4-8 μ long; conidiophores arising from the mycelium, scattered or crowded, 2-2.5 μ wide, phialides grouped in whorls or heads in the upper part of the conidiophores or occasionally from short lateral hyphae, narrowly fusoid-ovoid, acuminate above, 8-12 × 2 μ; conidia hyaline, fusoid-ellipsoid, 2-3 × 1.5-2 μ, catenulate.

On Opilionoidea, south fork of Kaukonahua, 1100 ft., Waianuka, Oahu, Hawaii, Dec. 9, 1945, D. P. Rogers, 2090, type. (Univ. of Mich. and N. Y. Bot. Gard.)

The hosts of this collection are so severely parasitized that accurate determination is difficult. They appear to be arachnids belonging to the Opilionoidea. The conidial stage resembles *Spi-*

¹ Paper from the Herbarium and the Department of Botany of the University of Michigan.

caria longipes Petch. In the latter species the phialides generally develop from short hyphal branches. Petch (7) states that *S. longipes* is the conidial stage of *Torrubiella gonyleptica* (Möller) Petch. *T. gonyleptica* differs from *T. pulvinata* in having the




FIG. 1. *Torrubiella pulvinata* and *T. confragosa*.

UNIVERSITY OF MICHIGAN LIBRARY

pe
de
de
sh
sir
pr

I
ver
pat
mu
br
ten
for
cyl
sur
mu
I
Ba

ex
ver
sup
350
wa
bro
4 μ
for
not
dri
int
shc
nat
pre

(
Br.
Ba
Pla
I
rul

species the phialides generally
 es. Petch (7) states that *S.*
rubiella gonylepticida (Möller)
 on *T. pulvinata* in having the



rubiella and *T. confragosa*.

perithecia scattered over the entire body and appendages of the host developing from a very scanty superficial mycelium. Möller (2) describes the perithecia of *T. gonylepticida* as orange red, flask-shaped, 300–400 μ long, and the asci as $170 \times 3 \mu$. Since it is desirable to have a name for the conidial stage, *Spicaria pulvinata* is proposed for that of *T. pulvinata*.

***Torrubiella confragosa* sp. nov. (FIG. 1, C)**

Mycelium tenue, album vel cremeum, hospitem tegente, floccosum, pulverulentum; perithecia irregulariter sparsa vel congesta, superficialia vel paululum immersa, ovoidea, $350-650 \times 200-375 \mu$, rubro-brunnea vel castanea, muris $20-35 \mu$ crassis ex duobus stratis constructis, interno strato rubro-brunneo, externo strato brunneo-flavo; asci cylindrici, $200-350 \times 3.5-4 \mu$, tenuibus membranis, ad apices membranis spissatis $2-2.5 \mu$; ascospores filiformes, multiseptatae; synnemata summe brevia, $300-800 \mu$ longa, 80μ crassa, cylindrica, interdum furcata; phialides $14-20 \mu$ longae, deorsum 1.5μ latae, sursum attenuatae, acuminatis apicibus; conidia oblonga, $3-4 \times 1.5-1.7 \mu$, mucositate tecta, 2–5 in glebulis congregata.

In Coccidis, Novo Petropolis, Brazil, May 1923, Rick, specimen typicum; Bayeaux, Haiti, J. R. Weir.

Mycelium thin, white to cream color, covering scale insects and extending slightly beyond on the substratum, slightly tufted, pulverulent; perithecia irregularly scattered to crowded over the scale, superficial or slightly embedded at the base in the mycelium, ovoid, $350-650 \times 200-375 \mu$, reddish brown to dark chestnut brown, the wall $20-35 \mu$ thick and consisting of two layers, the inner reddish brown, the outer brownish yellow; asci cylindric, $200-350 \times 3.5-4 \mu$, the wall thin, thickened at the apex, $2-2.5 \mu$; ascospores filiform, 0.5μ wide, almost as long as asci, multiseptate, part-spores not seen; synnemata very short, $300-800 \mu$ long, 80μ thick, cylindrical or furcate with short irregular branches, consisting of loosely interwoven hyphae; phialides arising from the outer hyphae or from short lateral hyphae, $14-20 \mu$ long, 1.5μ wide below, gradually narrowing to an acuminate apex; conidia oblong, $3-4 \times 1.5-1.7 \mu$, produced in a mucus, often in clumps of 2 to 5.

On large scale-insects. Novo Petropolis, Rio Grande do Sud, Brazil, May 1923, Rick, type (Univ. Mich. and Farlow Herb.); Bayeaux, Haiti, J. R. Weir (Univ. Mich. and Mycol. Coll. Bur. Plant Ind. Washington, D.C.).

Both of these collections were received as *Torrubiella rubra*. *T. rubra* develops perithecia only at the margin of the mycelium cover-

ing the scale or on a hypothallus and the asci are much larger than those of *T. confragosa*. Petch (3) describes a conidial stage for *T. rubra* which differs considerably from that of *T. confragosa*. The latter can best be classified as a *Hirsutella*, although the synnemata are poorly developed. The scattered phialides and clumped spores in a surrounding mucus are characteristic of that genus. Since the conidial stage of fungi of this group frequently develop without the perithecial stage it is convenient to have a name applying to it and consequently *Hirsutella confragosa* is proposed.

Torrubiella liberiana sp. nov. (FIG. 2, A)

Perithecia albedo-brunnea, ex myceliis in articulis hospitis erumpentia, partim immersa, ovoidea, 300-400 × 150-200 μ; asci cylindrici, tennes, 210 μ longa, 3-4 μ crassa, tenuibus membranibus, ad apices membranibus spissatis, 2 μ; ascospores filiformes, multiseptatae, fragentes, segmentis 3-4 × 0.5-0.7 μ.

Ex formica, Bonata, Liberia, Dec. 8, 1947, J. T. Baldwin, Jr.

Perithecia light brown, developing in small white patches of mycelium at the joints of the appendages and of the body, partly embedded, ovoid, 300-400 × 150-200 μ; asci cylindric, slender, up to 210 μ long, 3-4 μ wide, wall thin, thickened at the apex, 2 μ; ascospores filiform, nearly as long as the asci, multiseptate, breaking into one-celled fragments, 3-4 × 0.5-0.7 μ.

On an ant. Bonata, Central Province, Liberia, Dec. 8, 1947, J. T. Baldwin, Jr. (Univ. Mich.), **type**.

As far as the writer is aware, this is the first report of a species of *Torrubiella* on ants. It occurs in close association with synnemata of a *Hirsutella* which may be the conidial stage. However, it was separated from a collection of ants which were mostly parasitized by *Cordyceps australis*. A *Gibellula* also occurs on a few specimens. The relationship of these fungi is therefore very uncertain and it seems best to describe them separately.

Hirsutella liberiana sp. nov. (FIG. 2, B)

Synnemata pauca, sparsa, ex exiguis albidis myceliis plerumque in articulis erumpentia, albida, tenuia, cylindrica, 3 mm. longa, deorsum 100 μ crassa, sursum paululum attenuata, simplicia vel interdum furcata; phialides sparsae vel congregatae, plerumque nonnullae ex brevibus lateralibus hyphis erumpentes, deorsum fusoside-ellipsoideae vel fusoside-oblongae, 8-12 × 3 μ,



FIG. 2.

is and the asci are much larger than 1 (3) describes a conidial stage for erably from that of *T. confragosa*. d as a *Hirsutella*, although the syn- ed. The scattered phialides and ng mucus are characteristic of that ge of fungi of this group frequently ul stage it is convenient to have a equently *Hirsutella confragosa* is

na sp. nov. (FIG. 2, A)

myceliis in articulis hospitis erumpentia, < 150-200 μ ; asci cylindrici, tenues, 210 μ ranis, ad apices membranis spissatis, 2 μ ; , fragentes, segmentis 3-4 \times 0.5-0.7 μ . e. 8, 1947, J. T. Baldwin, Jr.

veloping in small white patches of appendages and of the body, partly 50-200 μ ; asci cylindric, slender, up thin, thickened at the apex, 2 μ ; as- g as the asci, multiseptate, breaking \times 0.5-0.7 μ .

ul Province, Liberia, Dec. 8, 1947, ch.), **type**.

, this is the first report of a species occurs in close association with syn- ay be the conidial stage. However, on of ants which were mostly para-

A *Gibellula* also occurs on a few of these fungi is therefore very un- rible them separately.

ia sp. nov. (FIG. 2, B)

ignis albidis myceliis plerumque in arti- ylindrica, 3 mm. longa, deorsum 100 μ simplicia vel interdum furcata; phialides nonnullae ex brevibus lateralibus hyphis oideae vel fusoidae-oblongae, 8-12 \times 3 μ ,




FIG. 2. *Torrubiella*, *Hirsutella* and *Gibellula*

sursum attenuatae quisque 10-14 μ longo sterigmate; conidia cylindrica, 3-4 \times 0.7-1.0 μ ; muco tecta, in glebulas congregata.

Ex formica, Bonata, Liberia, Dec. 8, 1947, J. T. Baldwin, Jr.

Synnemata few, scattered, arising from a slight whitish mycelium, usually from the joints of the body and the appendages, whitish, slender, cylindric, slightly narrowing upward, up to 3 mm. long, 100 μ thick, simple or occasionally branched above; phialides scattered to crowded, usually several arising from short, one-celled lateral hyphal branches from the outer hyphae of the synnemata, the lower part fusoid-ellipsoid to fusoid-oblong, 8-12 \times 3 μ , the upper part prolonged into an attenuated, acuminate sterigma, 10-14 μ long, each producing several conidia which adhere together in a persistent mucus; conidia cylindric, rounded at both ends, 3-4 \times 0.7-1.0 μ .

On an ant, Bonata, Central Province, Liberia, Dec. 8, 1947, J. T. Baldwin, Jr. (Univ. Mich.), **type**.

Petch (6) has described *Hirsutella formicarum* on ants from British Guiana and Ceylon and has stated that it is the conidial stage of *Cordyceps unilateralis*. He describes the conidia as narrowly cymbiform, 9-11 \times 2 μ . In an earlier publication (4) he described the conidia of the Ceylon specimens as oval, 3-5 \times 1 μ . Kobayasi (1) has identified Japanese collections on ants as *H. formicarum*, describing the conidia as ovoid, ellipsoid or fusiform, 3-4.2 \times 1.5-2 μ . It would seem probable that more than one species is concerned. *Hirsutella liberiana* is distinguished by its cylindric narrow conidia. It was associated with perithecia of *Torrubiella liberiana* which is described herein.

Hirsutella ramosa sp. nov. (FIG. 2, C)

Synnemata multa, gregaria, tenuia, irregulariter ramosa, 8 mm. longa, deorsum 52-140 μ crassa, sursum attenuata, acuminate apicibus, deorsum albido-flava, sursum alba, multis albis rectis vel obliquis ramulis; phialides sparsae vel 1-3 caespitosae, hyalinae, subulatae, 19.3-36.4 μ longae, deorsum 2.1-3.2 μ crassae, sursum attenuatae, acuminate apicibus; conidia oblonga, 3.2-5.5 \times 1.1-1.4 μ , tenui muco tecta, in parvas globosas glebulas 4.3-6.4 μ dia. congregata.

Ex larva lepidopteri, Salmon River, Nova Scotia, Sept. 7, 1931, L. E. Wehmeyer (1474).

Synnemata numerous, arising from all parts of the host, slender, irregularly branched, the main stem up to 8 mm. long, 52-140 μ

thick at the base, gradually yellow below, white at angles or slightly oblique secondary branches, tubular and parallel, minutely roughened, at the base, gradually narrower upper part of the stem arising singly directly nema or 1-3 at the apical cylindrical hyphae; conidia a slight mucus. adhering

On fragments of a lepidopteran, Salmon River, Nova Scotia, Sept. 7, 1931, L. E. Wehmeyer (Univ. Mich.).

The much branched synnemata distinguish *H. ramosa* from other *Hirsutella* Lepidoptera.

Gibellula form

Synnemata sparsa, plerumque brunnea, anguste cylindrica, hyphis laxis brunneis asperis inferioribus cellulis 3-5 μ longis, 4-8 μ lati, hyalina, globosis phialidium gerentibus; prophialides quisque paucam phialidem gerentes, 1.5-2.5 μ ; conidia hyalinae, 3-5 \times 1.5-2.5 μ ; conidia hyalinae. Ex formicis, Belleyella-K. Jr., specimen typicum; Bon.

Synnemata scattered, arising from all parts of the host, slender, irregularly branched, the main stem up to 8 mm. long, 52-140 μ thick at the base, gradually yellow below, white at angles or slightly oblique secondary branches, tubular and parallel, minutely roughened, at the base, gradually narrower upper part of the stem arising singly directly nema or 1-3 at the apical cylindrical hyphae; conidia a slight mucus. adhering

o sterigmate; conidia cylindrica, 3-4 gregata.

1947, J. T. Baldwin, Jr.

; from a slight whitish mycelium, ly and the appendages, whitish, ing upward, up to 3 mm. long, dly branched above; phialides al arising from short, one-celled ter hyphae of the synnemata, the usoid-oblong, 8-12 x 3 μ, the uated, acuminate sterigma, 10- onidia which adhere together in ic, rounded at both ends, 3-4 x

ince, Liberia, Dec. 8, 1947, J. T.

tella formicarum on ants from stated that it is the conidial stage scribes the conidia as narrowly lier publication (4) he described s as oval, 3-5 x 1 μ. Kobayasi ons on ants as *H. formicarum*, ipsoid or fusiform, 3-4.2 x 1.5- more than one species is con- stinguished by its cylindrical nar- with perithecia of *Torrubiella*

o. nov. (FIG. 2, C)

irregulariter ramosa, 8 mm. longa, uata, acuminate apicibus, deorsum ectis vel obliquis ramulis; phialides bulatae, 19.3-36.4 μ longae, deorsum uminate apicibus; conidia oblonga, i parvas globosas glebulas 4.3-6.4 μ

Nova Scotia, Sept. 7, 1931, L. E.

om all parts of the host, slender, m up to 8 mm. long, 52-140 μ

thick at the base, gradually narrowing to acuminate apices, light yellow below, white above, the branches numerous, white, at right angles or slightly oblique to the main stem, occasionally producing secondary branches, the hyphae in the stem and branches longitudinal and parallel, multiseptate, 2 μ wide; phialides hyaline, minutely roughened, subulate, 19.3-36.4 μ long, 2.1-3.2 μ wide at the base, gradually narrowing to an acuminate apex, scattered on the upper part of the stem and branches giving a setose appearance, arising singly directly from a cell of the outer hypha of the synnema or 1-3 at the apex of short, 6.4-10.7 μ, laterally projecting cylindrical hyphae; conidia oblong, 3.2-5.5 x 1.1-1.4 μ, covered with a slight mucus, adhering to form small, 4.3-6.4 μ, spherical clumps.

On fragments of a lepidopterous larva on a mossy log, Salmon River, Nova Scotia, Sept. 7, 1931, L. E. Wehmeyer, 1474, type (Univ. Mich.).

The much branched synnemata and narrow subulate phialides distinguish *H. ramosa* from other species of *Hirsutella* infecting Lepidoptera.

Gibellula formicarum sp. nov. (FIG. 2, D & E)

Synnemata sparsa, plerumque ex articulis hospitis erumpentia, albido-brunnea, anguste cylindrica, 1-2 mm. longa, 50-150 μ crassa, exterioribus hyphis laxis brunneis asperulatis; conidiophora 50-150 μ longa, 2-4-septata, inferioribus cellulis 3-5 μ latis, asperulatis, superiore cella obovoidea, 7-8 x 4-8 μ, levi, hyalina, globosum vel cylindricum capitulum prophialidium et phialidium gerentis; prophialides ellipsoideae vel subglobosae, 3.5-4 x 2.5-3 μ, quisque paucam phialidem gerentis; phialides ovoideae vel cylindricae, 4-14 x 1.5-2.5 μ; conidia hyalina, cylindrica, 3-4.5 x 1-1.5 μ.

Ex formicis, Belleyella-Kondessu-Zui, Liberia, Dec. 1947, J. T. Baldwin, Jr., specimen typicum; Bonata, Liberia, Dec. 8, 1947, J. T. Baldwin, Jr.

Synnemata scattered, arising from various parts of the body and appendages of the host, usually at the joints, very light brown, narrowly cylindrical, 1-2 mm. long, 50-150 μ thick, composed of longitudinal somewhat interwoven hyphae, the outer, loose, brown, asperulate; conidiophores arising from short lateral prolongations of cells of the outer hyphae, 50-150 μ long, 2-4-septate, the lower cells 3-5 μ wide, the walls brownish, asperulate, the terminal cell obovoid, 7-8 x 4-8 μ, smooth, hyaline, bearing a globose to cylindrical head of prophialides and phialides; prophialides ellipsoid to subspherical, 3.5-4 x 2.5-3 μ, each bearing several phialides; phialides ovoid to cylindrical, 4-14 x 1.5-2.5 μ; conidia hyaline, cylindrical, rounded at the ends, 3-4.5 x 1-1.5 μ.

On ants, along the route Belleyella-Kondessu-Zui, Western Province, Liberia, Dec. 8, 1947, J. T. Baldwin, Jr. (Univ. Mich.).

The specimens were included in collections of *Cordyceps australis* (Speg.) Sacc. which were received from Dr. Baldwin. The collections were composites collected from several localities. The synnemata of *Gibellula formicarum* were not found associated with clavae of *C. australis* upon the same insect and therefore the ascogenous stage is uncertain. Petch (5) has reported that the conidial stage of *C. australis* is an *Hymenostilbe*.

LITERATURE CITED

1. Kobayasi, Y. The genus *Cordyceps* and its allies. Tokyo Bun. Dar. Sci. Rep. 5 (no. 84): 53-260. 1941.
2. Möller, Alfred. Phycomyceten und Ascomyceten. In Schimper's Bot. Mit. Tropen. Heft 9: 319 pp. 1901.
3. Petch, T. Studies in entomogenous fungi. III. *Torrubiella*. Trans. British Myc. Soc. 9: 108-128. 1924.
4. ——. Studies in entomogenous fungi. *Ibid.* 10: 28-45. 1924.
5. ——. Notes on entomogenous fungi. *Ibid.* 16: 209-245. 1931.
6. ——. Notes on entomogenous fungi. *Ibid.* 19: 161-194. 1934.
7. ——. Notes on entomogenous fungi. *Ibid.* 21: 34-67. 1938.

EXPLANATION OF FIGURES

FIG. 1. A & B. *Torrubiella pulvinata*. A. Pulvinate mass of mycelium bearing perithecia arising from a leg of the host. $\times 12$. B. Body of a host covered by a mass of mycelium and conidiophores. $\times 7$. C. *Torrubiella confragosa*. Large scale-insects covered with a thin layer of mycelium-bearing synnemata and perithecia. $\times 4$.

FIG. 2. A. *Torrubiella liberiana* showing perithecia developing on a small mass of mycelia at the base of an antenna of an ant. $\times 25$. B. *Hirsutella liberiana* showing synnemata arising from a small patch of mycelium on an ant. $\times 7$. C. *Hirsutella ramosa* showing the branched synnemata. $\times 7$. D & E. *Gibellula formicarum*. D. An ant showing the scattered synnemata on various parts of the body. $\times 7$. E. A portion of a synnema showing capitate conidiophores (stained with nigrosin). $\times 200$.